

# Current Perspectives in the Diagnosis and Treatment of Nontuberculous Mycobacterial Pulmonary Disease

Michael R Loebinger<sup>1</sup> and Tobias Welte<sup>2</sup>

Consultant Respiratory Medicine, Royal Brompton Hospital, London, UK; 2. Department of Respiratory Medicine, Hannover Medical School, Hannover, Germany

DOI: <http://doi.org/10.17925/ERPD.2016.02.02.54>

**N**ontuberculous mycobacteria (NTM), an extremely diverse group with a wide spectrum of virulence, appear to be rising in prevalence. This may be due to both increased awareness and improved diagnosis, among other factors. Most NTM-pulmonary disease (NTM-PD) infections are caused by the slow growing *Mycobacterium avium complex* (MAC) however, other species can also cause pulmonary disease including *M. abscessus*, *M. xenopi*, *M. malmoense* and *M. kansasii*. Symptoms of NTM-PD vary considerably and may include chronic cough, with or without sputum production or haemoptysis. Treatment options are limited at present and associated with significant intolerance, treatment failures and drug–drug interactions. Further prospective studies are needed to define suitable drug regimens for different NTM species and to improve knowledge of the natural history and epidemiology of NTM-PD.

## Keywords

Nontuberculous mycobacteria (NTM), *Mycobacterium avium complex* (MAC), *Mycobacterium avium*, *Mycobacterium intracellulare*, epidemiology, diagnosis, treatment

**Disclosure:** Michael R Loebinger has received consultancy fees from Insmmed, but this has not impacted on the article. Tobias Welte has received consultancy fees from Insmmed, but this has not impacted on the article.

**Acknowledgement:** Medical writing assistance was provided by Catherine Amey at Touch Medical Media, London and funded by Insmmed.

**Open Access:** This article is published under the Creative Commons Attribution Noncommercial License, which permits any non-commercial use, distribution, adaptation and reproduction provided the original author(s) and source are given appropriate credit.

**Received:** 9 June 2016

**Accepted:** 25 July 2016

**Citation:** *European Respiratory & Pulmonary Diseases*, 2016;2(2):54–7

**Corresponding Author:** Michael R Loebinger, Respiratory Medicine, Royal Brompton Hospital, Sydney Street, London, SW3 6NP, UK. E: M.Loebinger@rbht.nhs.uk

**Support:** The publication of this article was supported by Insmmed. The views and opinions expressed are those of the authors and not necessarily those of Insmmed.

Nontuberculous mycobacteria (NTM) are a diverse group of bacteria with a wide spectrum of virulence.<sup>1</sup> They are a microaerobic, non-motile organisms that have a lipid-rich, hydrophobic cell wall that is substantially thicker compared with most other bacteria.<sup>2</sup> NTM are associated with biofilm formation, which contributes to disinfectant- and antibiotic-resistance.<sup>3–5</sup> Many of these organisms are also resistant to high temperature and low pH.<sup>6–8</sup> Ubiquitous in the environment, NTM are found with the greatest concentrations in water sources, both natural and treated, and in soil sources.<sup>9</sup> There are currently over 170 species of mycobacteria and more are likely to be discovered in the future owing to improved culturing techniques and molecular technology advances.<sup>10</sup>

The majority of NTM-pulmonary disease (NTM-PD) infections are caused by the slow growing *Mycobacterium avium complex* (MAC) that includes *Mycobacterium avium* and *Mycobacterium intracellulare*, although other species including *M. abscessus*, *M. xenopi*, *M. malmoense* and *M. kansasii* can also cause lung disease. Not all NTM are considered pathogenic, with species such as *Mycobacterium gordonae* commonly isolated but not usually associated with the development of NTM-PD and a wide variability reported in the clinical relevance of species.<sup>12</sup> Co-infection with different NTM strains or other bacteria has also been reported.<sup>9,13</sup> NTM lung disease is characterised by clinical, microbiological and radiological criteria.<sup>1</sup> High resolution computed tomography (CT) scanning may consist of a variety of disease patterns including nodular/bronchiectatic disease, fibrocavitary disease, solitary pulmonary nodule, disseminated disease, and hypersensitivity-like disease.<sup>1</sup> Acquisition of NTM is believed to be from the environment; however, emerging literature, using sequencing of NTM isolates, suggest that indirect cross infection of *M. abscessus* is possible in patients with cystic fibrosis.<sup>14</sup> This has led to changes in infection control standards in these patient groups.<sup>15,16</sup>

NTM are treated with antimicrobial chemotherapy, however treatment response is often poor, and side effects significant. Deciding which mycobacteria and patients to treat, and when, and the assessment of response to treatment is very difficult. The purpose of this review is to provide a succinct summary of issues surrounding the epidemiology, diagnosis and treatment of NTM-PD.

## Epidemiology

In contrast to tuberculosis, NTM isolation in pulmonary specimens does not necessarily equate with active pulmonary disease, thus the incidence and prevalence of NTM pulmonary infections is difficult to characterise accurately.<sup>9</sup> In addition, unlike tuberculosis, isolating NTM does not constitute a reportable condition in many countries. Population-based data for prevalence are available mainly from the US, Europe, New Zealand and Australia (where NTM isolation is reportable).<sup>17</sup> These estimates are based on a mixture of microbiology laboratory reports, medical

insurance returns and surveys. Annual prevalence estimates in North America and Australia range from 3.2 to 9.8 per 100,000.<sup>17</sup> In Europe the estimates are generally lower, up to 3.3/100,000.<sup>17,18</sup> The prevalence is significantly higher in certain regions and patient groups and period prevalence has been put at over 100/100,000 in the over 65-year-olds in the US over an 11-year period.<sup>19</sup> The incidence of NTM disease is increasing throughout the world,<sup>11,17</sup> with some estimates of an increase of 8.2% per year.<sup>19</sup> NTM-PD is currently more common in the US than tuberculosis.<sup>20</sup> Improvements in awareness and diagnosis may in part underlie this rise.

In addition to the difference in prevalence, the predominant NTM species has also been reported to differ substantially by geographical region.<sup>21</sup> To obtain information of NTM species distribution, in an NTM-Network European Trials Group (NET) collaborative study, species identification data for 20,182 patients, from 62 laboratories in 30 countries across six continents was collected.<sup>22</sup> In total, 91 different species were isolated; *M. avium* complex predominated in most areas, although *M. Kansaii* predominated in Poland and Slovakia and *M. Xenopi* predominated in Hungary. MAC was most frequently isolated in Japan, Australia and South Africa.<sup>19,23-25</sup>

Pre-existing chronic lung disease (bronchiectasis, cystic fibrosis, chronic obstructive pulmonary disease [COPD], idiopathic pulmonary fibrosis) predisposes patients to acquiring NTM respiratory infection.<sup>26,27</sup> Other disease risk factors include thoracic skeletal abnormalities, taking immunomodulatory drugs and rheumatoid arthritis.<sup>17</sup> Clustering of disease within families may suggest a heritable genetic predisposition to susceptibility of NTM-PD. Defects in interleukin-12 and interferon-gamma defence pathways may increase susceptibility to NTM infections<sup>28</sup> and these defects have been described both in individuals and in familial clustering of disease.<sup>29</sup> A recent study utilising exome sequencing of patients, family members and controls has suggested a multi-hit, multi-genetic predisposition to NTM disease.<sup>30</sup> In addition to host predisposition, environmental exposure is likely to play a significant role in predisposition to NTM-PD, as suggested by variance in prevalence and mortality between and within countries. In studies in the US,<sup>19</sup> Japan<sup>31</sup> and Australia,<sup>32</sup> temperature, humidity and alteration in soil components have impacted on NTM prevalence.

Mortality in NTM-PD is significant, with 5-year mortality after definite NTM disease at 40.1%, according to Danish population data.<sup>33</sup> In this study, patients with *M. xenopi* had the worst prognosis. A randomised controlled study of NTM treatment regimens by the British Thoracic Society demonstrated similarly high 5-year all-cause mortality rates of 38.8% for MAC, 48.5% for *M. malmoense* and 38.2% for *M. xenopi*.<sup>34</sup> Other retrospective studies have specifically looked at MAC lung disease and assessed specific characteristics associated with a worse prognosis. The 5-year all-cause mortality of patients with definite pulmonary MAC was 25.6% in a retrospective study (n=164) of patients with pulmonary MAC disease.<sup>35,36</sup> High Charlson comorbidity index scores and the presence of cavitary lesions were associated with higher mortality. Prognostic factors of all-cause mortality were identified in a retrospective analysis of 634 HIV-negative patients with MAC lung disease.<sup>37</sup> The all-cause 5-year mortality was 23.9% and negative prognostic factors of all-cause mortality included: male sex, older age, presence of comorbidity, body mass index (BMI) <18.5 kg/m<sup>2</sup>, anaemia, hypoalbuminemia and erythrocyte sedimentation rate ≥50 mm/h. In a retrospective review of 72 patients with nodular bronchiectatic MAC lung disease, lower BMI, cavities and macrolide-resistance at initial visit seemed to be

**Table 1: Summary of the American Thoracic Society diagnostic criteria for pulmonary nontuberculous mycobacterial infection<sup>1</sup>**

Clinical
1. Pulmonary symptoms, nodular or cavitary opacities on chest radiograph, or a high-resolution computed tomographic scan that shows multifocal bronchiectasis with multiple small nodules; And
2. Appropriate exclusion of other diagnoses.
Microbiologic
1. Positive culture results from at least two separate expectorated sputum samples (If the results from the initial sputum samples are nondiagnostic, consider repeat sputum AFB smears and cultures); Or
2. Positive culture results from at least one bronchial wash or lavage; Or
3. Transbronchial or other lung biopsy with mycobacterial histopathological features (granulomatous inflammation or AFB) and positive culture for NTM or biopsy showing mycobacterial histopathological features (granulomatous inflammation or AFB) and one or more sputum or bronchial washings that are culture positive for NTM
4. Expert consultation should be obtained when NTM are recovered that are either infrequently encountered or that usually represent environmental contamination
5. Patients who are suspected of having NTM lung disease but who do not meet the diagnostic criteria should be followed until the diagnosis is firmly established or excluded
6. Making the diagnosis of NTM lung disease does not, per se, necessitate the institution of therapy, which is a decision based on potential risks and benefits of therapy for individual patients

Source: Adapted from Griffith DE, Aksamit T, Brown-Elliott BA, et al., 2007.<sup>1</sup> AFB = acid-fast bacillus; NTM = nontuberculous mycobacteria

associated with progression to deteriorating disease over 5 years.<sup>38</sup> In another retrospective analysis, which involved data from 52 adults with non cystic fibrosis bronchiectasis and coexisting MAC infection over a 5-year period, chronic pulmonary aspergillosis and nodules with cavitation and emphysema on high-resolution CT scans at presentation were identified as a strong independent predictors of mortality.<sup>39</sup> In a retrospective cohort study of 124 patients with NTM, predominately *M. avium*, the most important predictor for increased mortality was the radiological appearance of a consolidation.<sup>40</sup> Direct comparisons between studies are difficult due to the differences in treatment regimens, patient cohorts and methodologies.

## Diagnosis

NTM are ubiquitous in the environment and isolation from sputum does not always signify disease. The American Thoracic Society (ATS) and the Infectious Disease Society of American (IDSA) jointly published guidelines on diagnosis of NTM infections in 2007 (Table 1).<sup>1</sup> An early differential diagnosis of NTM lung disease from pulmonary tuberculosis is important because the therapeutic regimen differs from that of pulmonary tuberculosis and it is not necessary to track the contacts of patients with NTM-PD.<sup>41</sup>

Symptoms are variable although chronic cough with or without sputum production or haemoptysis, is common. Other symptoms include fatigue, malaise, dyspnoea, fever, chest pain and weight loss. The evaluation of NTMs is often complicated by coexisting lung diseases including cystic fibrosis, bronchiectasis and pneumoconiosis. For patients in whom diagnosis is not clear, expert consultation should be sought. High resolution CT scans showing NTM-PD are presented in Figure 1.

**Figure 1: High resolution computed tomography scans revealing nontuberculous mycobacterial pulmonary disease**



A: Cavernous *Mycobacterium intracellulare* infection in a patient with non cystic fibrosis bronchiectasis, deterioration with standard treatment; B: Consolidated *Mycobacterium chimaera* (MAC group)-infection in a patient with non cystic fibrosis bronchiectasis; C: Tree-in-bud sign in a chronic obstructive pulmonary disease patient with *Mycobacterium avium* infection; D: Persistent *Mycobacterium intracellulare* infection patient with A1 antitrypsin deficiency; E: *Mycobacterium avium* infection in a patient with Lady Windermere syndrome, middle lobe destroyed.

## Treatment

The 2007 ATS/IDSA guidelines are the most commonly followed worldwide. They recommend that treatment for most patients with nodular/bronchiectatic MAC disease involves a three-times-weekly regimen of clarithromycin (1,000 mg) or azithromycin (500 mg), rifampin (600 mg) and ethambutol (25 mg/kg), although in many cases daily treatment may be appropriate.<sup>42</sup> For patients with fibrocavitary MAC lung disease or severe nodular/bronchiectatic disease, a daily regimen of clarithromycin (1,000 mg) or azithromycin (250 mg), rifampin (600 mg) or rifabutin (150–300 mg) and ethambutol (15 mg/kg) with consideration of three-times-weekly amikacin or streptomycin early in therapy is recommended. Patients should be treated until culture negative for 1 year while receiving antibacterial therapy.<sup>1</sup> Similar combinations of drugs have also been suggested for other NTM that commonly cause NTM-PD such as *M. kansasii*, *M. malmoense* and *M. xenopi*.<sup>1</sup> No established regimens of proven efficacy exist for *M. abscessus*; although standard treatment based on case series consists of an induction phase of intravenous therapy including medications such as amikacin, carbapenem, tigecycline or ceftazidime, followed by a continuation phase of oral antibiotics.<sup>43</sup> A macrolide-based regimen is often used and surgical debridement may also be important. However, macrolide resistance is common in *M. abscessus* because the organism possesses an inducible macrolide-resistance gene.<sup>44,45</sup> In some instances more than one NTM species is present and there are no data available to guide the clinician in these cases so they need to be evaluated and treated on an individual basis.<sup>13</sup>

In some patients, clinical improvement and culture conversion will be the aim, however, for others, stability will be a more realistic target.<sup>1</sup> The potential toxicity of anti-mycobacterial therapy must be

**Table 2: Typical adverse events requiring monitoring that are associated with drugs used to treat NTM infections<sup>46</sup>**

Drug	Adverse events
Clarithromycin	Gastrointestinal, QT prolongation
Azithromycin	Gastrointestinal, QT prolongation
Amikacin	Vestibular/auditory, renal,
Rifampin	Hepatotoxicity, flu-like syndrome
Ciprofloxacin	Gastrointestinal, tendonitis, QT prolongation
Levofloxacin	Gastrointestinal, tendonitis, QT prolongation
Moxifloxacin	Gastrointestinal, tendonitis, QT prolongation
Ethambutol	Optic neuritis
Linezolid	Neuropathy, thrombocytopenia, myelosuppression, optic neuritis
Isoniazid	Liver, peripheral neuropathy
Tigecycline	Gastrointestinal
Cefoxitin	Gastrointestinal, seizures
Imipenem	Gastrointestinal, seizures

Adapted from Egelund EF, Fennelly KP, Peloquin CA, 2015.<sup>46</sup>

**Table 3: Typical drug–drug interactions involving nontuberculous mycobacteria**

Nontuberculous mycobacteria treatment	Important drug–drug interactions	Reference
Macrolides	Affect oxidation of many drugs via interaction with CYP, including: alprazolam, midazolam, clozapine, carbamazepine, simvastatin, lovastatin, warfarin and cyclosporine. May interact with direct thrombin inhibitors and factor Xa inhibitors. In particular, clarithromycin increases the anticoagulant effect of rivaroxaban and dabigatran	Westphal (2000) <sup>50</sup> Delavenne et al., (2013) <sup>51</sup>
Rifamycin drugs (rifampin, rifabutin and rifapentine)	Strong inducer of CYP	Baciewicz (2013) <sup>52</sup>
Fluoroquinolones	Inhibitors of CYP. Inhibit metabolism of NSAIDs and theophylline	Davey (1988) <sup>53</sup>

CYP = Cytochrome P-450; NSAIDs = nonsteroidal anti-inflammatory drugs.

balanced against a sometimes indolent clinical course. A patient's comorbidities and quality-of-life issues are also important to take into account. There are currently insufficient data on the natural history of untreated infections to guide clinical decision-making in this respect. In clinical practice it is often difficult to discern whether or not disease progression is due to NTM. Even if NTM is diagnosed, lung disease treatment may not always be needed. For patients with indolent NTM lung disease there is usually enough time for careful consideration of the optimal management. Once a risk-benefit decision has been made to begin NTM therapy, the choice of agents and duration of treatment is based on the specific organism and extent of disease. Before a decision is made to treat with the aim of improving symptoms such as cough, it should be considered whether the symptoms are attributable to a concomitant diagnoses of bronchiectasis, sinus disease and/or gastroesophageal reflux disease.



## Limitations of current treatment

The drugs used for treatment of NTM have not been developed specifically for these infections but rather, the rationale for their use has often been extrapolated from the treatment of tuberculosis and are often based on personal experiences of the treating clinician. The long duration of treatment often leads to adverse drug reactions, increased potential for drug–drug interactions and nonadherence to treatment or treatment discontinuation.<sup>46</sup> Typical adverse events of drugs used to treat NTM infections and drug–drug interactions are summarised in *Tables 2* and *3*. NTM infections are common in elderly people and age-related changes in drug absorption, metabolism and excretion may lead to a decline in efficacy and increased toxicity.<sup>47</sup>

## Treatment practices

Adherence to the 2007 ATS/IDSA guidelines for the treatment of NTM-PD in the US has been revealed to be poor, with suboptimal antibiotic regimens commonly prescribed across physician specialities.<sup>48</sup> Overall, 95% of the 1,286 physicians selected were in the following specialities: pulmonology, infectious diseases (IDs), internal medicine or family/general practice. Out of 349 physicians who completed questionnaires,

only 13% of antibiotic regimens prescribed for patients with MAC, were in accordance to ATS/IDSA guidelines. Around half (56%) did not include a macrolide and 16% were for macrolide monotherapy.<sup>48</sup>

## Conclusions and future perspectives

NTM-PD is increasing in prevalence, leading to an emerging public health problem.<sup>17,49</sup> More work is needed to investigate species-specific environmental niches as causes of differences in global isolation patterns.<sup>22</sup> Better management of the disease remains an unmet need with improvements required in diagnosis, treatment, patient adherence to treatment and patient outcome. Present treatment options are limited, with drug regimens associated with significant failure and intolerance. Individualised treatment should be administered after careful consideration of both the efficacy and risk of adverse events.

New treatments for NTM infections are imperative and more prospective studies are warranted to define suitable drug regimens for different NTM species and to better define the natural history and epidemiology of this condition. Due to the complexity of the problem, involvement of specialist centres is important. □

- Griffith DE, Aksamit T, Brown-Elliott BA, et al., An official ATS/IDSA statement: diagnosis, treatment, and prevention of nontuberculous mycobacterial diseases, *Am J Respir Crit Care Med*, 2007;175:367–416.
- James J Plorde, Mycobacteria. In: KJ Ryan, CG Ray, Sherris Medical Microbiology, 4th Edition: McGraw-Hill Medical, 2004;439–56.
- Jarlier V, Nikaïdo H, Mycobacterial cell wall: structure and role in natural resistance to antibiotics, *FEMS Microbiol Lett*, 1994;123:11–8.
- Schulze-Robbeke R, Janning B, Fischeder R, Occurrence of mycobacteria in biofilm samples, *Tubercle and Lung Disease*, 1992;73:141–4.
- Falkinham JO 3rd, Growth in catheter biofilms and antibiotic resistance of *Mycobacterium avium*, *J Med Microbiol*, 2007;56:250–4.
- Bodmer T, Miltner E, Bermudez LE, *Mycobacterium avium* resists exposure to the acidic conditions of the stomach, *FEMS Microbiol Lett*, 2000;182:45–9.
- Kirschner RA, Jr., Parker BC, Falkinham JO 3rd, Epidemiology of infection by nontuberculous mycobacteria. *Mycobacterium avium*, *Mycobacterium intracellulare*, and *Mycobacterium scrofulaceum* in acid, brown-water swamps of the southeastern United States and their association with environmental variables, *Am Rev Respir Dis*, 1992;145:271–5.
- Faria S, Joao I, Jordao L, General overview on nontuberculous mycobacteria, biofilms, and human infection, *J Pathog*, 2015;2015:809014.
- Johnson MM, Odell JA, Nontuberculous mycobacterial pulmonary infections, *J Thorac Dis*, 2014;6:210–20.
- Tortoli E, Impact of genotypic studies on mycobacterial taxonomy: the new mycobacteria of the 1990s, *Clin Microbiol Rev*, 2003;16:319–54.
- Prevots DR, Shaw PA, Strickland D, et al., Nontuberculous mycobacterial lung disease prevalence at four integrated health care delivery systems, *Am J Respir Crit Care Med*, 2010;182:970–6.
- van Ingen J, Bendien SA, de Lange WC, et al., Clinical relevance of non-tuberculous mycobacteria isolated in the Nijmegen-Arnhem region, The Netherlands, *Thorax*, 2009;64:502–6.
- Cowman S, Burns K, Benson S, et al., The antimicrobial susceptibility of non-tuberculous mycobacteria, *J Infect*, 2016;72:324–31.
- Bryant JM, Grogono DM, Greaves D, et al., Whole-genome sequencing to identify transmission of *Mycobacterium abscessus* between patients with cystic fibrosis: a retrospective cohort study, *Lancet*, 2013;381:1551–60.
- Cystic Fibrosis Trust, Cross-infection, 2016. Available at: [www.cysticfibrosis.org.uk/life-with-cystic-fibrosis/cross-infection](http://www.cysticfibrosis.org.uk/life-with-cystic-fibrosis/cross-infection) (accessed date: 26 July 2016).
- Floto RA, Olivier KN, Saiman L, et al., US Cystic Fibrosis Foundation and European Cystic Fibrosis Society consensus recommendations for the management of non-tuberculous mycobacteria in individuals with cystic fibrosis: executive summary, *Thorax*, 2016;71:88–90.
- Prevots DR, Marras TK, Epidemiology of human pulmonary infection with nontuberculous mycobacteria: a review, *Clin Chest Med*, 2015;36:13–34.
- Ringshausen FC, Wagner D, de Roux A, et al., Prevalence of nontuberculous mycobacterial pulmonary disease, Germany, 2009–2014, *Emerg Infect Dis*, 2016;22:1102–5.
- Adjemian J, Olivier KN, Seitz AE, et al., Prevalence of nontuberculous mycobacterial lung disease in U.S. Medicare beneficiaries, *Am J Respir Crit Care Med*, 2012;185:881–6.
- Iseman MD, Marras TK, The importance of nontuberculous mycobacterial lung disease, *Am J Respir Crit Care Med*, 2008;178:999–1000.
- Marras TK, Daley CL, Epidemiology of human pulmonary infection with nontuberculous mycobacteria, *Clin Chest Med*, 2002;23:553–67.
- Hoefsloot W, van Ingen J, Andrejak C, et al., The geographic diversity of nontuberculous mycobacteria isolated from pulmonary samples: an NTM-NET collaborative study, *Eur Respir J*, 2013;42:1604–13.
- Adjemian J, Olivier KN, Seitz AE, et al., Spatial clusters of nontuberculous mycobacterial lung disease in the United States, *Am J Respir Crit Care Med*, 2012;186:553–8.
- Feazel LM, Baumgartner LK, Peterson KL, et al., Opportunistic pathogens enriched in showerhead biofilms, *Proc Natl Acad Sci U S A*, 2009;106:16393–9.
- Thomson RM, Changing epidemiology of pulmonary nontuberculous mycobacteria infections, *Emerg Infect Dis*, 2010;16:1576–83.
- Aksamit TR, Phillely JV, Griffith DE, Nontuberculous mycobacterial (NTM) lung disease: the top ten essentials, *Respir Med*, 2014;108:417–25.
- Winthrop KL, Chang E, Yamashita S, et al., Nontuberculous mycobacteria infections and anti-tumor necrosis factor- $\alpha$  therapy, *Emerg Infect Dis*, 2009;15:1556–61.
- Dorman SE, Holland SM, Interferon- $\gamma$  and interleukin-12 pathway defects and human disease, *Cytokine Growth Factor Rev*, 2000;11:321–33.
- Field SK, Fisher D, Cowie RL, *Mycobacterium avium* complex pulmonary disease in patients without HIV infection, *Chest*, 2004;126:566–81.
- Szymanski EP, Leung JM, Fowler CJ, et al., Pulmonary nontuberculous mycobacterial infection. A multisystem, multigenic disease, *Am J Respir Crit Care Med*, 2015;192:618–28.
- Morimoto K, Iwai K, Ohmori M, et al., [Nontuberculous mycobacteriosis mortality in Japan] Article in Japanese, *Kekkaku*, 2011;86:547–52.
- Chou MP, Clements AC, Thomson RM, A spatial epidemiological analysis of nontuberculous mycobacterial infections in Queensland, Australia, *BMC Infect Dis*, 2014;14:279.
- Andrejak C, Thomsen VQ, Johansen IS, et al., Nontuberculous pulmonary mycobacteriosis in Denmark: incidence and prognostic factors, *Am J Respir Crit Care Med*, 2010;181:514–21.
- Jenkins PA, Campbell IA, Banks J, et al., Clarithromycin vs ciprofloxacin as adjuncts to rifampicin and ethambutol in treating opportunistic mycobacterial lung diseases and an assessment of *Mycobacterium vaccae* immunotherapy, *Thorax*, 2008;63:627–34.
- Ito Y, Hirai T, Maekawa K, et al., Predictors of 5-year mortality in pulmonary *Mycobacterium avium*-intracellulare complex disease, *Int J Tuberc Lung Dis*, 2012;16:408–14.
- Winthrop KL, McNeilley E, Kendall B, et al., Pulmonary nontuberculous mycobacterial disease prevalence and clinical features: an emerging public health disease, *Am J Respir Crit Care Med*, 2010;182:977–82.
- Hayashi M, Takayanagi N, Kanauchi T, et al., Prognostic factors of 634 HIV-negative patients with *Mycobacterium avium* complex lung disease, *Am J Respir Crit Care Med*, 2012;185:575–83.
- Kitada S, Uenami T, Yoshimura K, et al., Long-term radiographic outcome of nodular bronchiectatic *Mycobacterium avium* complex pulmonary disease, *Int J Tuberc Lung Dis*, 2012;16:660–4.
- Zoumot Z, Boutou AK, Gill SS, et al., *Mycobacterium avium* complex infection in non-cystic fibrosis bronchiectasis, *Respirology*, 2014;19:714–22.
- Gommans EP, Even P, Linssen CF, et al., Risk factors for mortality in patients with pulmonary infections with non-tuberculous mycobacteria: a retrospective cohort study, *Respir Med*, 2015;109:137–45.
- Koh WJ, Yu CM, Suh GY, et al., Pulmonary TB and NTM lung disease: comparison of characteristics in patients with AFB smear-positive sputum, *Int J Tuberc Lung Dis*, 2006;10:1001–7.
- Schonfeld N, Haas W, Richter E, et al., Recommendations for diagnosis and treatment of nontuberculous mycobacterioses of the German Central Committee against tuberculosis and the German Respiratory Society, *Pneumologie*, 2013;67:605–33.
- Jeon K, Kwon OJ, Lee NY, et al., Antibiotic treatment of *Mycobacterium abscessus* lung disease: a retrospective analysis of 65 patients, *Am J Respir Crit Care Med*, 2009;180:896–902.
- Griffith DE, The talking *Mycobacterium abscessus* blues, *Clin Infectious Dis*, 2011;52:572–4.
- Nash KA, Brown-Elliott BA, Wallace RJ Jr. A novel gene, *erm(41)*, confers inducible macrolide resistance to clinical isolates of *Mycobacterium abscessus* but is absent from *Mycobacterium chelonae*, *Antimicrob Agents Chemother*, 2009;53:1367–76.
- Egelund EF, Fennelly KP, Peloquin CA, Medications and monitoring in nontuberculous mycobacteria infections, *Clin Chest Med*, 2015;36:55–66.
- Mangoni AA, Jackson SH, Age-related changes in pharmacokinetics and pharmacodynamics: basic principles and practical applications, *Br J Clin Pharmacol*, 2004;57:6–14.
- Adjemian J, Prevots DR, Gallagher J, et al., Lack of adherence to evidence-based treatment guidelines for nontuberculous mycobacterial lung disease, *Ann Am Thorac Soc*, 2014;11:9–16.
- Kendall BA, Winthrop KL, Update on the epidemiology of pulmonary nontuberculous mycobacterial infections, *Semin Respir Crit Care Med*, 2013;34:87–94.
- Westphal JF, Macrolide - induced clinically relevant drug interactions with cytochrome P-450A (CYP) 3A4: an update focused on clarithromycin, azithromycin and dirithromycin, *Br J Clin Pharmacol*, 2000;50:285–95.
- Delavenne X, Ollier E, Basset T, et al., A semi-mechanistic absorption model to evaluate drug-drug interaction with dabigatran: application with clarithromycin, *Br J Clin Pharmacol*, 2013;76:107–13.
- Baciewicz AM, Chrisman CR, Finch CK, Self TH, Update on rifampin, rifabutin, and rifapentine drug interactions, *Curr Med Res Opin*, 2013;29:1–12.
- Davey PG, Overview of drug interactions with the quinolones, *J Antimicrob Chemother*, 1988;22 Suppl C:97–107.