# Advances in Pulmonary Care in Duchenne Muscular Dystrophy

Oscar H Mayer,<sup>1</sup> Erik K Henricson,<sup>2</sup> Craig M McDonald<sup>2</sup> and Gunnar M Buyse,<sup>3</sup> on behalf of the Cooperative International Neuromuscular Research Group (CINRG) and the DELOS study group

1. Division of Pulmonology, Children's Hospital of Philadelphia, Philadelphia, US; 2. Department of Physical Medicine and Rehabilitation, School of Medicine, University of California, Sacramento, US; 3. Paediatric Neurology, University Hospitals Leuven, Belgium

DOI: https://doi.org/10.17925/USN.2017.13.01.35

During the source of pulmonary function results in ineffective airway clearance, sleep-disordered breathing and nocturnal and daytime respiratory failure. Routine measures of pulmonary function include forced vital capacity (FVC) and peak expiratory flow (PEF). Both measures follow parallel trajectories and relentlessly decline, reaching the lower limit of normal of 80% of predicted at early teenage years. Moreover, decline in PEF and FVC are closely correlated with respiratory complications and clinically relevant thresholds for FVC are defined in standard of care recommendations. Glucocorticoids (GCs) delay the onset of pulmonary function decline, but once patients have reached the 80% of predicted threshold the decline of FVC and PEF in GC users and patients not using GCs is comparable. In the successful phase III DELOS trial in DMD patients not using GCs, the short-chain benzoquinone idebenone (Raxone®, Santhera Pharmaceuticals, Liestal, Switzerland) has demonstrated statistically significant and clinically relevant efficacy on expiratory and inspiratory function in patients in the pulmonary function decline stage. These results indicate that idebenone can modify the natural course of respiratory disease progression, which is relevant in clinical practice where loss of respiratory function continues to be a predominant cause of early morbidity and mortality in DMD.

#### Keywords

Duchenne muscular dystrophy, glucocorticoids, idebenone, pulmonary function, peak expiratory flow, forced vital capacity

**Disclosure**: All authors act as scientific consultants and advisors to Santhera Pharmaceuticals (Switzerland). Oscar H Mayer has additional consulting relationships with Bristol-Myers Squib, Marathon Pharmaceuticals, Catabasis Pharmaceuticals, Sarepta Pharmaceuticals, Fibrogen, Biogen, AveXis and Hoffman-LaRoche, and is an investigator in the SIDEROS trial. Erik K Henricson is study co-chair of the CINRG Duchenne natural history study and has served as a scientific consultant for Bristol Myers Squibb, PTC Therapeutics and Genzyme, Inc. He received travel assistance from Parent Project Muscular Dystrophy (USA). Craig M McDonald is study co-chair of the CINRG Duchenne natural history study and has served as a consultant for PTC, Prosensa, Sarepta, Eli Lilly, Pfizer, Halo Therapeutics, Cardero and Mitokyne, and serves on external advisory boards related to DMD for PTC and Eli Lilly. Gunnar M Buyse was investigator of clinical trials in Duchenne muscular dystrophy sponsored by Santhera Pharmaceuticals, Prosensa and GlaxoSmithKline, and is senior clinical investigator of the Research Foundation Flanders (FWO Vlaanderen, Belgium). He is also the inventor of relevant patent applications.

Acknowledgements: Medical writing assistance was provided by Katrina Mountfort at Touch Medical Media, London and funded by Santhera Pharmaceuticals. The authors thank Mika Leinonen (Santhera Pharmaceuticals, Switzerland) for statistical analyses and Thomas Meier (Santhera Pharmaceuticals, Switzerland) for contributing to the manuscript and supporting the preparation of analyses, figures and tables. Santhera Pharmaceuticals is the sponsor of the DELPHI, DELOS and SIDEROS trials and supported this publication.

Investigators of the Cooperative International Neuromuscular Research Group (CINRG): V Vishwanathan, S Chidambaranathan, W Douglas Biggar, Laura C McAdam, Jean K Mah, Mar Tulinius, Avital Cnaan, Lauren P Morgenroth, Robert Leshner, Carolina Tesi-Rocha, Mathula Thangarajh, Tina Duong, Andrew Kornberg, Monique Ryan, Yoram Nevo, Alberto Dubrovsky, Paula R Clemens, Hoda Abdel-Hamid, Anne M Connolly, Alan Pestronk, Jean Teasley, Tulio E Bertorini, Richard Webster, Hanna Kolski, Nancy Kuntz, Sherilyn Driscoll, John B Bodensteiner, Jose Carlo, Ksenija Gorni, Timothy Lotze, John W Day, Peter Karachunski, Erik K Henricson, Richard T Abresch and Craig M McDonald (for updated affiliations, see online version at www.ncbi.nlm.nih.gov/ pubmed/28139640).

Investigators of the DELOS study group (sorted by country): Austria: G Bernert, F Knipp (Vienna). Belgium: GM Buyse, N Goemans, M Van den Hauwe (Leuven). France: T Voit, V Doppler, T Gidaro (Paris); J-M Cuisset, S Coopman (Lille). Germany: U Schara, S Lutz (Essen); J Kirschner, S Borell, M Will (Freiburg). Italy: MG D'Angelo, E Brighina, S Gandossini (Lecco); K Gorni, E Falcier (Milan); L Politano, P D'Ambrosio, A Taglia (Naples). The Netherlands: JJGM Verschuuren, CSM Straathof (Leiden). Spain: JJ Vilchez Padilla, N Muelas Gómez (Valencia). Sweden: T Sejersen, M Hovmöller (Stockholm). Switzerland: P-Y Jeannet, C Bloetzer (Lausanne). USA: S Iannaccone, D Castro (Dallas); G Tennekoon, R Finkel, C Bönnemann (Philadelphia); C McDonald, E Henricson, N Joyce (Sacramento); S Apkon, RC Richardson (Seattle).

Authorship: All named authors meet the International Committee of Medical Journal Editors (ICMJE) criteria for authorship of this manuscript, take responsibility for the integrity of the work as a whole, and have given final approval to the version to be published.

Open Access: This article is published under the Creative Commons Attribution Noncommercial License, which permits any noncommercial use, distribution, adaptation, and reproduction provided the original author(s) and source are given appropriate credit.

Received: February 7, 2017

Accepted: March 13, 2017

Citation: US Neurology, 2017;13(1):35-41

Corresponding Author: Oscar H Mayer, Division of Pulmonology, The Children's Hospital of Philadelphia, 11309 Colket Center, 3510 Civic Center Boulevard, Philadelphia, PA 19104, US. E: MAYERO@email.chop.edu

Support: The publication of this article was supported by Santhera Pharmaceuticals. The views and opinions expressed are those of the authors and do not necessarily reflect those of Santhera Pharmaceuticals.



## Figure 1: Co-linear decline in FVC%p and PEF%p in patients with Duchenne muscular dystrophy

Data is mean  $\pm$  SEM (n=334); dotted lines at age 10 and 20 years: period of approximately linear decline in PEF%p and FVC%p. FVC%p = forced vital capacity percent predicted; PEF%p = peak expiratory flow percent predicted; PFT = pulmonary function test. Source: CINRG Duchnne natural history study

Duchenne muscular dystrophy (DMD), the most common and devastating type of muscular dystrophy,<sup>1</sup> is characterised by the absence of the protein dystrophin, which causes premature muscle cell failure and leads to progressive muscle atrophy and loss. The condition is typically diagnosed at age 3-5 years when children start to show signs of physical disability including difficulty in walking. Patients experience progressive muscle weakness and become non-ambulant in early teenage years.<sup>2</sup> From the age of 10-12 years, pulmonary function typically starts to decline<sup>3-5</sup> and cardio-pulmonary complications are the major cause of morbidity and early mortality in DMD.<sup>6</sup> Progressive pulmonary insufficiency requires the use of non-invasive and, in some circumstances, invasive ventilation, which, following the loss of ambulation, is the second irreversible disease milestone in DMD greatly impacting the patients' quality of life. Although standard of care recommendations, including recommendations to monitor pulmonary function, have been proposed, 2,6-8 overall compliance with these recommendations generally appears to be poor in real-world DMD care.9

DMD patients and their caregivers consider the preservation of pulmonary function, particularly maintaining effective cough and reducing the risk of airway infections, an important treatment goal,<sup>10</sup> which at the current time represents a significant unmet need.

The aim of this review is to provide an update on advances in the understanding of DMD, its natural history and recent pharmacotherapies. We will summarize advances in the understanding of pulmonary function assessment in patients with DMD and the natural course of pulmonary function decline. We will also report clinically relevant thresholds of pulmonary morbidity in the context of standard of care recommendations. Finally, we will review the effects of glucocorticoids (GCs) on pulmonary function outcomes and present emerging evidence that the investigational drug idebenone can slow the loss of pulmonary function in DMD patients.

# Assessment of pulmonary function in patients with Duchenne muscular dystrophy

Serial assessment of pulmonary function is a critical element of routine monitoring for patients with DMD, as it may enable early identification

and treatment of pulmonary complications. In the absence of obstructive pulmonary disease, forced vital capacity (FVC, measured in litres), peak expiratory flow (PEF, measured in litres/minute), and forced expiratory volume in 1 second (FEV1, measured in litres) are well-established dynamic spirometry measures useful in the assessment of restrictive pulmonary function changes due to neuromuscular weakness. In DMD, both inspiratory and expiratory respiratory muscle weakness can be indicated by low peak inspiratory and expiratory flows.<sup>5,11</sup> As pulmonary function is influenced by body growth and age, these measures are typically normalised relative to height and age12,13 and expressed as a percent of predicted (PEF%p, FVC%p, FEV1%p). As assessment of standing height in patients with DMD could be complicated by an inability to stand fully erect, scoliosis and joint contractures (particularly in non-ambulant patients), height can be estimated from ulna length measures.<sup>14,15</sup> Although effort dependent, FVC%p and PEF%p can reliably be measured in schoolage patients as shown by low within-subject coefficients of variation (CV) for successive measures.16

### Natural course of pulmonary function decline in Duchenne muscular dystrophy

Two recent prospective studies investigated the natural course of pulmonary function loss in DMD. The Cooperative International Neuromuscular Research Group (CINRG) Duchenne natural history study (DNHS) is an ongoing, prospective international natural history study and currently represents the most comprehensive description of a longitudinal observational cohort of DMD subjects.<sup>4,17</sup> Independently, prospective data in DMD patients were collected in the Neuromuscular Clinic at the Children's Hospital of Philadelphia from 2005–2010 as part of an Institutional Review Board approved United Dystrophinopathy Project (UDP) cohort study.<sup>5</sup>

As *Figure 1* from the CINRG-DNHS data set shows, flow-related PEF%p and volume-related FVC%p decline in a similar pattern, with decline in PEF%p slightly preceding the decline in FVC%p. At around 10–11 years of age, both parameters fall below the 80% predicted threshold, which is generally defined as the lower limit of normal and defines restrictive respiratory disease or low lung volume.<sup>7</sup> From the 80% threshold, both PEF%p and FVC%p follow a co-linear decline, approaching a floor of approximately 20% of predicted from age 20 onwards.

This parallel and tightly correlated decline between FVC%p and PEF%p from early teenage years was independently reported from the data set collected at the Neuromuscular Clinic at the Children's Hospital of Philadelphia.<sup>5</sup>

From this and other data collections, yearly rates of absolute decline in FVC%p and PEF%p can be extracted (*Table 1*). Although not directly comparable due to differences in age, GC-use status and other factors at time of assessment, these data indicate a yearly decline well above 5% absolute for both pulmonary function test (PFT) measures.

### Correlation between pulmonary function and general disease stage in Duchenne muscular dystrophy

It would be of clinical relevance if loss of pulmonary function could be predicted from or at least correlated with the general disease stage of DMD. As seen from *Figure 1*, crossing the 80% of predicted threshold for PEF%p or FVC%p and subsequent decline generally coincides with the time DMD patients become non-ambulant during early teenage years.

Reference	Patients (n)	Age (y)	Yearly rate of decline (absolute)		Comments
			FVC%p	PEF%p	
16, 18	33	15.0 ± 2.5	8.7%	8.9%	All patients GC non-users (>1 year prior to Baseline); Baseline: all patients with PEF%p <80%; mean Brook score 4.5; 93.9% non-ambulatory
19, 20	47	17.5 ± 4.7	4.68%		All patients non-ambulatory and GC non-users
5	60	5–24	5.0%	5.8%	Mix of GC-users and GC non-users
21	48 (28)	6–19	4.9%		28 patients with successive assessments; all patients GC non-users
4, 22	107*	10–18*	4.38%-5.85%¶	4.10%-6.78%	*for patients 10–18 years; majority of patients are GC users;¶ range for age brackets 10–12,13–15, 16–18 y
23	194	np	8.83% (Group A)* 7.52% (Group B)* 6.03% (Group C)*		All patients GC non-users; *groups divided by age at loss of ambulation (see text for details)

#### Table 1: Reports of contemporaneous natural history reference data for yearly rate of decline in PEF%p and FVC%p

Data are mean ± SD. FVC%p = forced vital capacity percent predicted; GC = glucocorticoids; np = not provided; PEF%p = peak expiratory flow percent predicted.

This is supported by baseline data from the DELOS trial<sup>18</sup> in 10–18-year-old DMD patients (n=64) where the majority of enrolled patients (92%) were already non-ambulant and at baseline had declined in pulmonary function to PEF%p of 53.8% (standard deviation [SD] 11.8) and FVC%p of 52.8% (SD 18.1).<sup>16,18</sup> The predictive value of the age of loss of ambulation was more systematically analysed in a French database with 278 DMD patients.<sup>23</sup> This study separated three groups of patients according to their age when they became non-ambulant (age at loss of ambulation – group A: before 8 years; group B: 8–<11 years; group C: 11–<16 years). Longitudinal analyses indicated that early loss of ambulation correlated with patients beginning to lose lung volume sooner, or having a peak vital capacity sooner (group A: 10.26 years; group B: 12.45 years; group C: 14.58 years).<sup>23</sup> Moreover, the annual rate of decline in FVC%p was also largest in patients who became non-ambulant at a very young age compared to patients who became non-ambulant at an older age (*Table 1*).

Earlier work by Anne Connolly and co-workers of the Muscular Dystrophy Association DMD Clinical Research Network<sup>19,20</sup> motivated attempts to correlate pulmonary function status with the patient's score on the Brooke Upper Extremity Functional Scale.<sup>24</sup> The Brooke scale scores upper extremity function from '1' (patient able to abduct arms into a full circle until they touch above the head), to '6' (hands cannot be raised to the mouth). For patients enrolled in the CINRG DNHS it could be shown that successive loss of upper extremity function is correlated with progressive loss of pulmonary function (Figure 2). Specifically, pulmonary function decline becomes apparent when patients reach Brooke score 2 or 3. which marks the transition of whether a patient can or cannot reach overhead. Of particular relevance is the transition from Brooke score 4 to score 5 (representing whether the patient still can raise the hand to the mouth or is losing this functional ability). A Brooke score of 5 appears to be an indicator that FVC%p has likely dropped below 50%, which is a clinically relevant threshold of pulmonary function loss (see below). These findings were independently confirmed by baseline data from the DELOS trial.<sup>18</sup> Here, patients with Brooke score 1–4 on average had PEF%p, FVC%p and FEV1%p of 60-80% of predicted, whereas in patients who lost the ability to raise their hand to their mouth (Brooke score 5), mean values for PEF%p, FVC%p and FEV1%p dropped below 50%.16

In summary, recent data indicate that pulmonary function loss can be correlated with the general disease stage of the patient. The age of loss of

# Figure 2: Declining FVC%p and Brooke Upper Extremity Function score in Duchenne muscular dystrophy



FVC%p (median, 75% and 25% quartiles and outliers) by Brooke's scale category (X-axis). Solid line: FVC%p of 80%. FVC%p = forced vital capacity percent predicted; OHR = overhead reach. Source: CINRG Duchenne natural history study

ambulation or the upper extremity function status may therefore provide readily available clinical assessments, which could indicate the need for more intense pulmonary function monitoring and clinical intervention.

# Sleep-disordered breathing in children and adolescents with Duchenne muscular dystrophy

In DMD, progressive respiratory muscle weakness will eventually cause nocturnal hypoventilation and later daytime hypoventilation and the need for continuous mechanical ventilation. However, nocturnal hypoventilation is typically considered more broadly within the category of sleep-disordered breathing (SDB). SDB is most likely in rapid eye movement (REM) sleep, a period characterized of maximal muscle atonia.<sup>25,26</sup> Clinical presentations of SDB can be restlessness in sleep, unrefreshing sleep or daytime fatigue.<sup>27</sup> In DMD patients, SDB is often aggravated by pharyngeal muscle weakness leading to obstructive sleep apnoea (OSA), typically caused by snoring.27 The most comprehensive study published to date by Sawnani and co-workers28 indicates that OSA during REM sleep is the commonest form of SDB in younger DMD patients. Interestingly, this work identified high body mass index (BMI) as a risk factor for OSA, which is of relevance due to the risk of weight gain during GC treatment. Identifying daytime predictors of SDB is still an area of active clinical research in DMD.27 For example, occurrence

## Table 2: Clinically relevant thresholds of FVC%p and standard of care recommendations

FVC%p category	Patient status	Standard of care recommendation	Reference
<80%	Established pulmonary function decline	Routine pulmonary function testing recommended (2x/year)	7
<50%	Mild to moderate pulmonary insufficiency	Special precautions during surgery (e.g. scoliosis surgery); recommended postoperative use of non-invasive ventilation	32
<40%	Signs and symptoms of hypoventilation	Volume recruitment / deep lung inflation techniques; manual and mechanical assisted cough techniques	6, 32
<30%	High risk of hypoventilation	Nocturnal ventilation; mandatory postoperative use of non-invasive ventilation	32

of hypopnoeas, nocturnal hypercapnic hypoventilation in REM sleep and continuous hypoventilation appeared to be correlated with a drop of inspiratory vital capacity.<sup>25,26</sup> Another study<sup>29</sup> established a drop in FEV1%p to <40% as a sensitive but not specific marker to predict sleep hypoventilation and hypoxic burden. Clearly, the identification of reliable and sensitive daytime predictors for SDB should constitute an area of future pulmonary function research in DMD.

# Characterization of pulmonary morbidity in Duchenne muscular dystrophy

Although FVC%p and PEF%p follow a parallel trajectory of decline, as shown in Figure 1, decline in FVC or FVC%p is historically the best validated pulmonary function predictor of early morbidity and mortality in patients with DMD.7,30 As soon as patients are non-ambulant and FVC%p drops below 80%, regular (2x/year) consultation by a physician specializing in paediatric respiratory care is recommended.<sup>7</sup> Once FVC%p drops below 50%, DMD patients may experience early clinical symptoms such as nocturnal hypoventilation, waking at night, morning headaches and problems with concentration and are considered at increased risk for post-procedural respiratory insufficiency when they undergo general anaesthesia or procedural sedation. The risk increases further once the FVC%p declines below 30%.31 Hence, the probabilities to present moderate pulmonary insufficiency is linked to FVC%p of <50% and severe pulmonary insufficiency in DMD is considered when patients have FVC%p <30%,623 and preoperative training in and postoperative use of noninvasive ventilation is recommended for patients with FVC%p <50% but is necessary for patients with FVC%p <30%.32 The risk of post-operative respiratory complications can be minimized with aggressive use of airway clearance and non-invasive ventilation as a bridge from invasive ventilation. It can be particularly helpful to introduce the assisted airway clearance strategies in the preoperative period if the patient does not already have an effective airway clearance regimen.

In addition, a drop in FVC%p below 40% is recommended as a threshold below which manual and mechanical assisted cough techniques are necessary to support effective airway clearance.<sup>4,32</sup> Transition between these clinically relevant pulmonary function categories commands a change in patient care and management, as summarised in *Table 2*.

# Figure 3: Annual rate of hospitalizations respiratory events by PEF%p and FVC%p category



(A) 247 patients with at least one PEF%p value of <80% were categorized according to their PEF%p value at any observation (number of observations, number of events) as follows; <80% to  $\geq$ 50% (204, 8); <50% to  $\geq$ 40% (89, 4); <40% to  $\geq$ 30% (88, 9); <30% (105, 43). (8) 230 patients with at least one FVC%p value of <80% were categorized according to their FVC%p value at any observation (number of observations, number of events) as follows; <80% to  $\geq$ 50% (201, 8); <50% to  $\geq$ 40% (74, 4); <40% to  $\geq$ 30% (67, 10); <30% (97, 45). FVC%p = forced vital capacity percent predicted; PEF%p = peak expiratory flow percent predicted. Source: CINRG Duchenne natural history study

Like a drop in FVC%p, a decline in PEF%p is also directly correlated with bronchopulmonary complications, such as airway infections, and the risk of hospitalization for any respiratory reason, as shown from data obtained at the CINRG DNHS (*Figure 3*).

Peak cough flow (PCF) is the peak flow generated during a cough and is used to objectively define airway clearance. Specifically, when the PCF falls below 160 l/min, the likelihood of successful extubation and airway clearance during an acute respiratory illness declines substantially, and patients with a PCF below 270 l/min are likely to fall below 160 l/min when viral illnesses occur and have problems with airway clearance.<sup>4,32–35</sup> However, this risk can be managed with the use of assisted airway clearance therapy. Therefore, once PCF declines below 270 l/min it is recommended that patients be started on assisted airway clearance with the cough assist device.<sup>32,33</sup> This is important because in the face of inadequate airway clearance, an otherwise benign upper respiratory tract infection can lead to lower respiratory tract infections and place the patient at risk of acute respiratory failure.

Clearly, the introduction of ventilatory support in routine clinical care for patients with DMD has considerably increased life expectancy<sup>36</sup> and recent reports indicate that median age of survival in patients using ventilator support is presently around 27 years or above.<sup>37,38</sup> Nevertheless, respiratory failure and associated complications are still a main cause for mortality even in ventilated patients<sup>39</sup> and in patients using GCs.<sup>40</sup>

### Glucocorticoids delay the onset but not the rate of pulmonary function loss in Duchenne muscular dystrophy

There is general agreement that ambulatory patients with DMD benefit from GC treatment,<sup>41</sup> often started already in patients as young as ~5 years of age.<sup>2</sup> The therapeutic objective of GC treatment is to slow the decline in muscle strength and stabilize pulmonary function. The use of GC has been shown to prolong ambulation by several years,<sup>42</sup> but the majority of GC-using patients become non-ambulant by approximately 14 years of age. More recently, continued treatment after the patient becomes nonambulant has also shown clinical benefit, such as reduction in the risk of progressive scoliosis, stabilization of pulmonary function and delay in the loss of upper limb function.<sup>4,43-47</sup> The results of these and other clinical reports support the use of GCs, which have become the mainstay of DMD disease management.<sup>48-50</sup> However, their chronic use in clinical practice has been hampered by significant adverse events and not all patients respond to GC to the same extent.

The effect of GC use on pulmonary function has been analysed from the CINRG DNHS.<sup>4,17</sup> These data show that onset of pulmonary function loss is delayed by ~2–3 years in patients receiving GCs compared to patients not on GC therapy (*Figure 4*). However, further analyses of the CINRG dataset show that once pulmonary function decline is established (i.e. falling below 80% of predicted), the year-to-year drop in mean FCV%p by year of age on average is similar in all patients (i.e. those using GCs vs. GC-naïve or previously GC-treated patients) down to 20 years of age where the FVC%p of GC-naïve patients approaches a nadir. Importantly, despite the use of GC, pulmonary function decline continues linearly and unabated from the age of approximately 10 years through to the early twenties (*Figure 4*).

This is important in light of the well-described risks associated with chronic administration of GCs such as growth retardation, bone demineralization and increased fracture risk, obesity, insulin resistance and hypertension, among others.<sup>2</sup> In the CINRG DNHS, 38% of DMD patients were either GC-naïve (24%) or had discontinued GC treatment (14%), with a clear trend to a higher proportion of GC non-users in the later post-ambulatory phase of the disease. In fact, for the patient group 10 years and older, 42% of DMD patients had never used GCs or discontinued their use because of side effects and tolerability limitations.<sup>4</sup> Interestingly, a study of all-cause mortality showed that the number of deaths due to respiratory failure was not significantly influenced by the GC use status of patients.<sup>40</sup>

### Therapeutic approach with idebenone to preserve pulmonary function in patients with Duchenne muscular dystrophy

There is an increasing body of literature indicating that dystrophin deficiency causes sarcolemmal fragility and intracellular Ca<sup>2+</sup> dysregulation, which generally leads to mitochondrial dysfunction, a reduction in resting ATP levels, an increased production of cell-damaging reactive oxygen species (ROS) and ultimately mitochondrial damage (reviewed in 51). Mitochondrial activation via stimulation of Complex II- or Complex III- mediated electron flux has been explored as a therapeutic approach to overcome mitochondrial dysfunction in dystrophin-deficient muscle.<sup>11</sup>

Idebenone, a synthetic short-chain benzoquinone, has been reported to utilize and activate Complex I-independent metabolic pathways<sup>52-54</sup> by the transfer of energy equivalents from the cytosol directly into the mitochondrial electron transport chain (ETC). Upon entering the cell, idebenone is efficiently reduced by the cytoplasmic enzyme NADH-quinone oxidoreductase 1 (NQO1) and the resulting reduced form of idebenone enters the mitochondria where it transfers its electrons to Complex III.<sup>52</sup> Preclinical studies showed that idebenone had a cardio-protective effect and improved exercise performance in the dystrophin-deficient mdx mouse model of DMD.<sup>55</sup>

A proof of concept, randomized, placebo-controlled trial (DELPHI trial) in 21 DMD patients treated for 12 months provided initial clinical

### Figure 4: Year-to-year drop in mean FVC%p over age classified by GC use status



Data is mean  $\pm$  SEM (n=334). FVC%p = forced vital capacity percent predicted. GC = glucocorticoid. Red arrows indicate shift in start of pulmonary function decline by QC use of about 2–3 years. Source: CINRG Duchenne natural history study

evidence that idebenone might have the potential to slow down loss of pulmonary function.  $^{\rm 56,57}$ 

Based on the outcome of the DELPHI trial, the confirmatory phase III, randomized, placebo-controlled DELOS trial was conducted, which investigated the efficacy of idebenone on pulmonary function outcomes. The trial enrolled 64 DMD patients (age 10–18 years, 92% non-ambulatory at baseline) who discontinued GC use at least 12 months prior to study start. Eligible patients had to be in the pulmonary function decline stage, defined as PEF%p at baseline of <80%. Patients received idebenone (Raxone<sup>®</sup> [Santhera Pharmaceuticals, Liestal, Switzerland], three times 300 mg/day with meals) or matching placebo for 52 weeks. The trial met its primary endpoint and demonstrated a statistically significant and clinically relevant reduction in the loss of PEF%p.<sup>18</sup>

The observed treatment difference (between idebenone and placebo groups) for PEF%p for the change during the one-year follow-up period in the intention to treat (ITT) population was 6.27% (95% confidence interval [CI]: 0.61–11.93; p=0.031). This magnitude for the observed treatment difference is comparable to the reported natural rate of decline over one year (see *Table 1*), thus the treatment difference is clinically relevant for patients not taking GCs. The observed treatment difference for FVC%p of 3.27% (95% CI: -0.43–6.97; p=0.082) corresponds to approximately half the magnitude of the expected decline over one year in untreated patients.

The outcome of the primary endpoint was robust across pre-specified study populations and supported by findings for changes in PEF, FVC%p, FVC, FEV1%p and FEV1 (reviewed in 11) and inspiratory function.<sup>58</sup> Interestingly, the treatment effect on PFTs was not influenced by the prior GC use status. Specifically, the estimated treatment effect for PEF%p and FVC%p was comparable in patients who never used GC and those who stopped GC use at least 12 month prior to study start.<sup>18</sup> In addition, idebenone showed a larger effect size for PEF%p and FVC%p in the subgroup of patients <14 years of age than in the older patients (>14 years), suggesting that patients may derive a larger benefit from idebenone if treatment is initiated early.<sup>18</sup>

#### Table 3: The idebenone group of the DELOS trial reported fewer bronchopulmonary events and hospitalizations due to respiratory causes compared to placebo

Analysis	Idebenone (n=31)	Placebo (n=33)	HR; (95% CI) p-value
Patients with at least one BAE	6	17	0.33; (0.13–0.83) <sup>1</sup> p=0.019
Total number of BAEs	7	28	0.28; (0.12–0.64) <sup>2</sup> p=0.003
Cumulative days with BAEs	82	222	
Patients hospitalized for respiratory reasons	1	4	
Total number of hospitalizations for respiratory reasons	1	6	0.20; (0.02–1.62) <sup>2</sup> p=0.130
Cumulative days in hospital	3	30	

BAEs counted: bronchitis, influenza (with respiratory symptoms), laryngitis, pneumonia, upper respiratory tract infection, viral infection (with respiratory symptoms), respiratory failure, acute respiratory failure, cough and dyspnoea. <sup>1</sup>Analysis is based on time to first event; <sup>2</sup>Analysis is based on time to all events, allowing multiple events per patient. BAEs = bronchopulmonary adverse events; CI = confidence interval; HR = hazard ratio.

Additional analyses of the DELOS trial showed that more patients in the placebo group compared to the idebenone group experienced bronchopulmonary adverse events (BAEs).59 In the placebo group there were 17 patients reporting 28 BAEs, compared to six patients reporting seven BAEs in the idebenone group. The hazard ratios (HR) calculated for the time to first BAE ('by patient') (HR 0.33, p=0.019) and for time to any BAE allowing multiple events per patient ('all BAEs') (HR 0.28, p=0.003) indicated a clear idebenone treatment effect (Table 3). The overall duration of BAEs was 222 days (placebo) versus 82 days (idebenone). The higher frequency of BAEs in the placebo group was also reflected in the number of serious adverse events leading to hospital admissions due to respiratory causes. There were four patients in the placebo group hospitalized six times for a total of 30 days, compared to only one patient in the idebenone group hospitalized once for three days (Table 3). Since mortality from respiratory failure in DMD occurs during acute respiratory illness requiring hospitalization, these findings of reduced respiratory illnesses and hospitalizations is an extremely valuable outcome.

In addition, there was also a difference in the use of systemic antibiotics utilized for the treatment of BAEs. In the placebo group, 13 patients (39.4%) reported 17 episodes of antibiotic use compared to seven patients (22.6%) reporting eight episodes of antibiotic use in the idebenone group. Furthermore, patients in the placebo group used systemic antibiotics for longer (105 days) compared to patients in the idebenone group (65 days).

# Table 4: Proportion of patients in whom FVC%p drops below clinically relevant threshold

Patients falling in FVC%p below	Idebenone (n=31) n (%)	Placebo (n=33) n (%)
50% predicted	4 of 20 (20%)	7 of 16 (44%)
40% predicted	4 of 24 (17%)	4 of 20 (20%)
30% predicted	4 of 29 (14%)	7 of 28 (25%)
Patients falling below any threshold*	10 of 29 (34%)	16 of 28 (57%)

\*Each patient is only counted once. FVC%p = forced vital capacity percent predicted.

As mentioned in *Table 2*, a drop in FVC%p below clinically relevant thresholds commands changes in patient care and management. When the data from the DELOS trial were evaluated relative to the FVC%p thresholds described earlier, there was also benefit with idebenone treatment. In the idebenone group at study entry, 20 patients presented with FVC%p above 50% and in four (20.0%), the FVC%p fell below 50% during the study. In the placebo group, at study entry there were 16 patients with FVC%p above 50% at baseline and seven (44%) declined below 50% during the study (*Table 4*). Similarly, higher proportions of patients in the placebo group had FVC%p below the 40% and 30% thresholds respectively.

As shown, there was a higher proportion of patients in the placebo group falling below any clinically relevant threshold for FVC%p (idebenone: 34%; placebo: 57%), resulting in a HR of 0.51 (95% CI: 0.23–1.14; p=0.099) in favour of the idebenone treatment group.

In summary, emerging evidence supports the therapeutic potential of idebenone in slowing the decline in pulmonary function in DMD patients. Current evidence for efficacy is available for patients in their teenage years presenting with advanced disease (majority of non-ambulant patients) who were already in the pulmonary function decline stage. Available data consistently demonstrate a treatment benefit with idebenone in patients who were not using GC treatment and for whom there is currently no alternative treatment option. Patients eligible for idebenone treatment could include patients who previously were treated with GCs or in whom GC treatment is not desired, not tolerated or is contraindicated.

A new phase III, randomized, placebo-controlled trial (SIDEROS trial) is currently ongoing with the objective to further assess the efficacy of idebenone in DMD patients by examining the impact on pulmonary function in the phase of decline (FVC%p <80%) and in patients receiving concomitant GC therapy.<sup>60</sup>

- Mah JK, Korngut L, Dykeman J, et al., A systematic review and meta-analysis on the epidemiology of Duchenne and Becker muscular dystrophy. *Neuromuscul Disord*, 2014;24:482–91.
   Bushby K, Finkel R, Birnkrant DJ, et al., Diagnosis and
- Bushby K, Finkel R, Birnkrant DJ, et al., Diagnosis and management of Duchenne muscular dystrophy, part 1: diagnosis, and pharmacological and psychosocial management, *Lancet Neurol*, 2010;9:77–93.
- Gayraud J, Ramonatxo M, Rivier F, et al., Ventilatory parameters and maximal respiratory pressure changes with age in Duchenne muscular dystrophy patients, *Pediatr Pulmonol*, 2010;45:552–9.
- Henricson EK, Abresch RT, Chaan A, et al., The cooperative international neuromuscular research group Duchenne natural history study: glucocorticoid treatment preserves clinically meaningful functional milestones and reduces rate of disease progression as measured by manual muscle testing and other commonly used clinical trial outcome measures, *Muscle Nerve*, 2013:48:55–67.
- 5. Mayer OH, Finkel RS, Rummey C, et al., Characterization of

pulmonary function in Duchenne Muscular Dystrophy, Pediatr Pulmonol, 2015;50:487–94.

- Bushby K, Finkel R, Birnkrant DJ, et al., Diagnosis and management of Duchenne muscular dystrophy, part 2: implementation of multidisciplinary care, *Lancet Neurol*, 2010;9:177–89.
- Finder JD, Birnkrant D, Carl J, et al., Respiratory care of the patient with Duchenne muscular dystrophy: ATS consensus statement, *Am J Respir Crit Care Med*, 2004;170:456–65.
- Birnkrant DJ, Ashwath ML, Noritz GH, et al., Cardiac and pulmonary function variability in Duchenne/Becker muscular dystrophy: an initial report. *J Child Neurol*. 2010;25:1110–5.
- Landfeldt E, Lindgren P, Bell CF, et al., Compliance to care guidelines for Duchenne muscular dystrophy. J Neuromuscul Dis, 2015;2:63–72.
- Hollin IL, Peay H, Apkon SD, et al., Patient-centered benefit-risk assessment in Duchenne muscular dystrophy, *Muscle Nerve*, 2016, doi: 10.1002/mus.25411. [Epub ahead of print]

- Buyse GM, Gueven N, McDonald C, Idebenone as a novel therapeutic approach for Duchenne muscular dystrophy, *European Neurological Review*. 2015;10:189–94.
- European Neurological Review, 2015;10:189–94.
   Quanjer PH, Stocks J, Polgar G, et al., Compilation of reference values for lung function measurements in children, Eur Respir J Suppl, 1989;4:1845–2615.
- Hankinson JL, Odencrantz JR, Fedan KB, Spirometric reference values from a sample of the general U.S. population, *Am J Respir Crit Care Med*. 1999:159:179–87.
- 14. Gauld LM, Kappers J, Carlin JB, et al., Height prediction from ulna length, *Dev Med Child Neurol*, 2004;46:475–80.
- Gauld LM, Kappers J, Carlin JB, et al., Prediction of childhood pulmonary function using ulna length, Am J Respir Crit Care Med, 2003;168:804–9.
- Meier T, Rummey C, Leinonen M, et al., Characterization of pulmonary function in 10–18 year old patients with Duchenne muscular dystrophy. *Neuromuscular Disorders*, 2017;27:307-314.
- McDonald CM, Henricson EK, Abresch RT, et al., The cooperative

international neuromuscular research group Duchenne natural history study—a longitudinal investigation in the era of glucocorticoid therapy: design of protocol and the methods used, Muscle Nerve. 2013;48:32–54.

- Buyse GM, Voit T, Schara U, et al., Efficacy of idebenone on respiratory function in patients with Duchenne muscular dystrophy not using glucocorticoids (DELOS): a doubleblind randomised placebo-controlled phase 3 trial, *Lancet*, 2015;385:1748–57.
- Comolly AM, Florence JM, Zaidman CM, et al., Clinical trial readiness in non-ambulatory boys and men with duchenne muscular dystrophy: MDA-DMD network follow-up, *Muscle Nerve*, 2016;54:681–9.
- Connolly AM, Malkus EC, Mendell JR, et al., Outcome reliability in non-ambulatory boys/men with Duchenne muscular dystrophy, *Muscle Nerve*, 2015;51:522–32.
- Khirani S, Ramirez A, Aubertin G, et al., Respiratory muscle decline in Duchenne muscular dystrophy, *Pediatr Pulmonol*, 2014;49:473–81.
- Abresch RT, McDonald CM, Henricson EK, et al., Pulmonary function characteristics of boys with Duchenne muscular dystrophy: data from the CINRG longitudinal study project, *Neuromuscul Disord*, 2013;23:738–852 (P.11.12).
   Humbertclaude V, Hamroun D, Bezzou K, et al., Motor and
- Humbertclaude V, Hamroun D, Bezzou K, et al., Motor and respiratory heterogeneity in Duchenne patients: implication for clinical trials, *Eur J Paediatr Neurol*, 2012;16:149–60.
   Brooke MH, Fenichel GM, Griggs RC, et al., Duchenne muscular
- Brooke MH, Fenichel GM, Griggs RC, et al., Duchenne muscular dystrophy: patterns of clinical progression and effects of supportive therapy. *Neurology*, 1989;39:475–81
- supportive therapy, *Neurology*, 1989;39:475–81.
  Mellies U, Ragette R, Schwake C, et al., Daytime predictors of sleep disordered breathing in children and adolescents with
- neuromuscular disorders, *Neuromuscul Disord*, 2003;13:123–8.
   Ragette R, Mellies U, Schwake C, et al., Patterns and predictors of sleep disordered breathing in primary myopathies, *Thorax*, 2002;57:724–8.
- Hoque R, Sleep-disordered breathing in Duchenne muscular dystrophy: an assessment of the literature, *J Clin Sleep Med*, 2016;12:905–11.
- Sawnani H, Thampratankul L, Szczesniak RD, et al., Sleep disordered breathing in young boys with Duchenne muscular dystrophy, J Pediatr, 2015;166:640-5 e1.
- Hukins CA, Hillman DR, Daytime predictors of sleep hypoventilation in Duchenne muscular dystrophy, Am J Respir Crit Care Med, 2000;161:166–70.
- Phillips MF, Quinlivan RC, Edwards RH, et al., Changes in spirometry over time as a prognostic marker in patients with Duchenne muscular dystrophy, *Am J Respir Crit Care Med*, 2001;164:2191–4.
- 31. Birnkrant DJ, Panitch HB, Benditt JO, et al., American College of Chest Physicians consensus statement on the respiratory

and related management of patients with Duchenne muscular dystrophy undergoing anesthesia or sedation, *Chest*, 2007;132:1977–84

- Birnkrant DJ, Bushby KM, Amin RS, et al., The respiratory management of patients with Duchenne muscular dystrophy: a DMD care considerations working group specialty article, *Pediatr Pulmonol*, 2010;45:739–48.
- Bach JR, Ishikawa Y, Kim H, Prevention of pulmonary morbidity for patients with Duchenne muscular dystrophy, *Chest*, 1997;112:1024–8
- Gauld LM, Boynton A, Relationship between peak cough flow and spirometry in Duchenne muscular dystrophy, *Pediatr Pulmonol*, 2005;39:457–60.
- Tzeng AC, Bach JR, Prevention of pulmonary morbidity for patients with neuromuscular disease, *Chest*, 2000;118:1390–6.
- Eagle M, Baudouin SV, Chandler C, et al., Survival in Duchenne muscular dystrophy: improvements in life expectancy since 1967 and the impact of home nocturnal ventilation, *Neuromuscul Disord*, 2002;12:926–9.
- Passamano L, Taglia A, Palladino A, et al., Improvement of survival in Duchenne muscular dystrophy: retrospective analysis of 835 patients, *Acta Myol*, 2012;31:121–5.
- Rall S, Grimm T, Survival in Duchenne muscular dystrophy, Acta Myol, 2012;31:117–20.
- Calvert LD, McKeever TM, Kinnear WJ, et al., Trends in survival from muscular dystrophy in England and Wales and impact on respiratory services, *Respir Med*, 2006;100:1058–63.
- Schram G, Fournier A, Leduc H, et al., All-cause mortality and cardiovascular outcomes with prophylactic steroid therapy in Duchenne muscular dystrophy, *JAm Coll Cardiol*, 2013;61:948–54.
- Matthews E, Brassington R, Kuntzer T, et al., Corticosteroids for the treatment of Duchenne muscular dystrophy, *Cochrane Database Syst Rev*, 2016;5:CD003725.
- Biggar WD, Gingras M, Fehlings DL, et al., Deflazacort treatment of Duchenne muscular dystrophy, *J Pediatr*, 2001;138:45–50.
- Biggar WD, Harris VA, Eliasoph L, et al., Long-term benefits of deflazacort treatment for boys with Duchenne muscular dystrophy in their second decade, *Neuromuscul Disord*, 2006;16:249–55.
- King WM, Ruttencutter R, Nagaraja HN, et al., Orthopedic outcomes of long-term daily corticosteroid treatment in Duchenne muscular dystrophy, *Neurology*, 2007;68:1607–13.
- Houde S, Filiatrault M, Fournier A, et al., Deflazacort use in Duchenne muscular dystrophy: an 8-year follow-up, *Pediatr Neurol*, 2008;38:200–6.
- Bello L, Gordish-Dressman H, Morgenroth LP, et al., Prednisone/ prednisolone and deflazacort regimens in the CINRG Duchenne Natural History Study, *Neurology*, 2015;85:1048–55.

- Pane M, Fanelli L, Mazzone ES, et al., Benefits of glucocorticoids in non-ambulant boys/men with Duchenne muscular dystrophy: a multicentric longitudinal study using the Performance of Upper Limb test, *Neuromuscul Disord*, 2015;25:749–53.
- Manzur AY, Kuntzer T, Pike M, et al., Glucocorticoid corticosteroids for Duchenne muscular dystrophy, *Cochrane Database Syst Rev*, 2008;1:CD003725.
- Bushby K, Lochmuller H, Lynn S, et al., Interventions for muscular dystrophy: molecular medicines entering the clinic, *Lancet*, 2009;374:1849–56.
- Gloss D, Moxley RT 3rd, Ashwal S, et al., Practice guideline update summary: corticosteroid treatment of Duchenne muscular dystrophy: report of the Guideline Development Subcommittee of the American Academy of Neurology. *Neurology*. 2016;86:465–72.
- Timpani CA, Hayes A, Rybalka E, Revisiting the dystrophin-ATP connection: how half a century of research still implicates mitochondrial dysfunction in Duchenne muscular dystrophy aetiology. *Med Hypotheses*, 2015;85:1021–33.
- Haefeli RH, Erb M, Gemperli AC, et al., NQ01-dependent redox cycling of idebenone: effects on cellular redox potential and energy levels, *PLoS One*, 2011;6:e17963.
- Giorgio V, Petronilli V, Ghelli A, et al., The effects of idebenone on mitochondrial bioenergetics, *Biochim Biophys Acta*, 2012;1817:363–9
- Erb M, Hoffmann-Enger B, Deppe H, et al., Features of idebenone and related short-chain quinones that rescue ATP levels under conditions of impaired mitochondrial complex I, *PLoS One*, 2012;7:e34153.
- Buyse GM, Van der Mieren G, Erb M, et al., Long-term blinded placebo-controlled study of SNT-MC17/idebenone in the dystrophin deficient mdx mouse: cardiac protection and improved exercise performance. *Fur Heart 1*, 2009;30:116–24.
- improved exercise performance, *Eur Heart J*, 2009;30:116–24.
   Buyse GM, Goemans N, van den Hauwe M, et al., Idebenone as a novel, therapeutic approach for Duchenne muscular dystrophy: results from a 12 month, double-blind, randomized placebocontrolled trial, *Neuromuscul Disord*, 2011;21:396–405.
- Buyse GM, Goemans N, Van den Hauwe M, et al., Effects of glucocorticoids and idebenone on respiratory function in patients with Duchenne muscular dystrophy, *Pediatr Pulmonol*, 2013;48:912–20.
- Buyse GM, Voit T, Schara U, et al., Treatment effect of idebenone on inspiratory function in patients with Duchenne muscular dystrophy. *Pediatr Pulmonol*, 2017;52:508-51.
   McDonald CM, Meier T, Voit T, et al., Idebenone reduces
- McDonald CM, Meier T, Voit T, et al., Idebenone reduces respiratory complications in patients with Duchenne muscular dystronby. *Neuromuscul Disord*. 2016;26:473–80.
- dystrophy, Neuromuscul Disord, 2016;26:473–80.
   A phase III double-blind study with idebenone in patients with Duchenne muscular dystrophy (DMD) taking glucocorticoid steroids (SIDEROS). Available at: https://clinicaltrials.gov/ct2/ show/NCT02814019 (Accessed 10/01/2017).