

**Personalized therapy for NSCLC:
Biomarker testing, treatment and
management in the presence of
MET alterations**

Disclaimer

- *Unapproved products or unapproved uses of approved products may be discussed by the faculty; these situations may reflect the approval status in one or more jurisdictions*
- *The presenting faculty have been advised by USF Health and touchIME to ensure that they disclose any such references made to unlabelled or unapproved use*
- *No endorsement by USF Health and touchIME of any unapproved products or unapproved uses is either made or implied by mention of these products or uses in USF Health and touchIME activities*
- *USF Health and touchIME accept no responsibility for errors or omissions*

A conversation between:



Dr Yasushi Goto

Assistant Chief
Division of Thoracic Oncology
National Cancer Center Hospital
Tokyo, Japan



Prof. Paul Paik

Associate Attending Physician
Memorial Sloan Kettering Cancer Center
New York, USA



Agenda

Evolving landscape of biomarker testing in advanced NSCLC

Identifying *MET*ex14 skipping mutations in NSCLC


Targeting *MET*ex14 skipping mutations in advanced NSCLC

METex14 skipping mutations in advanced NSCLC


Recommendations for biomarker testing^{1,2}

Biomarkers	NCCN guidelines	Pan-Asian guidelines
EGFR	✓	✓
ALK	✓	✓
ROS1	✓	✓
BRAF	✓	✓
PD-L1	✓	✓
METex14	✓	✗
NTRK1/2/3	✓	✗
KRAS	✓	✗
RET	✓	✗
ERBB2/HER2	✓	✗

Prevalence of oncogene mutations in NSCLC³



Mechanism of action of MET inhibitors^{4,5}



*EGFR mutations are prevalent in 10–16% of patients with NSCLC in Western populations and 40–50% in Asian populations.³

ALK, anaplastic lymphoma kinase; BRAF, v-raf murine sarcoma viral oncogene homolog B; EGFR, epidermal growth factor receptor; ERBB2, erb-b2 receptor tyrosine kinase 2; ex14, exon 14; HER2, human epidermal growth factor receptor 2; KRAS, Kirsten rat sarcoma; MET, mesenchymal-epithelial transition; NCCN, National Comprehensive Cancer Network; NSCLC, non-small cell lung cancer; NTRK, neurotrophic receptor tyrosine kinase; PD-L1, programmed death-ligand 1; RET, rearranged during transfection; ROS1, c-ros oncogene 1.


1. National Comprehensive Cancer Network (NCCN Guidelines®). Version 3.2023. Available at: https://www.nccn.org/professionals/physician_gls/pdf/nscl.pdf (accessed 17 April 2023); 2. Wu YL, et al. *Ann Oncol.* 2019;30:171–210; 3. Fois SS, et al. *Int J Mol Sci.* 2021;22:612; 4. Wu YL, et al. *Cancer Treat Rev.* 2021;95:102173; 5. Hong L, et al. *Ther Adv Med Oncol.* 2021;13:1758835921992976.

METex14 skipping mutations in advanced NSCLC


Recommendations for biomarker testing^{1,2}

Biomarkers	NCCN guidelines	Pan-Asian guidelines
EGFR	✓	✓
ALK	✓	✓
ROS1	✓	✓
BRAF	✓	✓
PD-L1	✓	✓
METex14	✓	✗
NTRK1/2/3	✓	✗
KRAS	✓	✗
RET	✓	✗
ERBB2/HER2	✓	✗

Prevalence of oncogene mutations in NSCLC³



Mechanism of action of MET inhibitors^{4,5}



*EGFR mutations are prevalent in 10–16% of patients with NSCLC in Western populations and 40–50% in Asian populations.³

ALK, anaplastic lymphoma kinase; BRAF, v-raf murine sarcoma viral oncogene homolog B; EGFR, epidermal growth factor receptor; ERBB2, erb-b2 receptor tyrosine kinase 2; ex14, exon 14; HER2, human epidermal growth factor receptor 2; KRAS, Kirsten rat sarcoma; MET, mesenchymal-epithelial transition; NCCN, National Comprehensive Cancer Network; NSCLC, non-small cell lung cancer; NTRK, neurotrophic receptor tyrosine kinase; PD-L1, programmed death-ligand 1; RET, rearranged during transfection; ROS1, c-ros oncogene 1.


1. National Comprehensive Cancer Network (NCCN Guidelines®). Version 3.2023. Available at: https://www.nccn.org/professionals/physician_gls/pdf/nscl.pdf (accessed 17 April 2023); 2. Wu YL, et al. *Ann Oncol.* 2019;30:171–210; 3. Fois SS, et al. *Int J Mol Sci.* 2021;22:612; 4. Wu YL, et al. *Cancer Treat Rev.* 2021;95:102173; 5. Hong L, et al. *Ther Adv Med Oncol.* 2021;13:1758835921992976.

METex14 skipping mutations in advanced NSCLC


Recommendations for biomarker testing^{1,2}

Biomarkers	NCCN guidelines	Pan-Asian guidelines
EGFR	✓	✓
ALK	✓	✓
ROS1	✓	✓
BRAF	✓	✓
PD-L1	✓	✓
METex14	✓	✗
NTRK1/2/3	✓	✗
KRAS	✓	✗
RET	✓	✗
ERBB2/HER2	✓	✗

Prevalence of oncogene mutations in NSCLC³



Mechanism of action of MET inhibitors^{4,5}



*EGFR mutations are prevalent in 10–16% of patients with NSCLC in Western populations and 40–50% in Asian populations.³

ALK, anaplastic lymphoma kinase; BRAF, v-raf murine sarcoma viral oncogene homolog B; EGFR, epidermal growth factor receptor; ERBB2, erb-b2 receptor tyrosine kinase 2; ex14, exon 14; HER2, human epidermal growth factor receptor 2; KRAS, Kirsten rat sarcoma; MET, mesenchymal-epithelial transition; NCCN, National Comprehensive Cancer Network; NSCLC, non-small cell lung cancer; NTRK, neurotrophic receptor tyrosine kinase; PD-L1, programmed death-ligand 1; RET, rearranged during transfection; ROS1, c-ros oncogene 1.

1. National Comprehensive Cancer Network (NCCN Guidelines®). Version 3.2023. Available at: https://www.nccn.org/professionals/physician_gls/pdf/nscl.pdf (accessed 17 April 2023); 2. Wu YL, et al. *Ann Oncol.* 2019;30:171–210; 3. Fois SS, et al. *Int J Mol Sci.* 2021;22:612; 4. Wu YL, et al. *Cancer Treat Rev.* 2021;95:102173; 5. Hong L, et al. *Ther Adv Med Oncol.* 2021;13:1758835921992976.