

The Excessive Carbon Footprint of Inhalers Used in Airway Disease and its Remedies

Laurence Désy, Philippe Lachapelle and Simon Couillard

Faculté de médecine et des sciences de la santé, Université de Sherbrooke, Sherbrooke, QC, Canada

The carbon footprint of human activities is a conversation topic worldwide. Many fields, including healthcare, have attempted to reduce their carbon footprint. For example, in Canada, 4.6% of greenhouse gases are manufactured by healthcare institutions. When it comes to the accountable parties in healthcare, the current treatments for asthma and chronic obstructive pulmonary disease are the main contributors. These highly prevalent diseases affect 11% and 10% of the Western populations, respectively. This editorial aims to discuss the impact of pressurized metered dose inhalers on the environment, alternatives and changes to current diagnostic and therapeutic practices, public awareness, potential changes to regulations and paths forward.

Keywords

Asthma, carbon dioxide, carbon footprint, chronic obstructive pulmonary disease, dry powder inhalers, global warming, inhalers, pressurized metered dose inhalers, vaporizers and nebulizers

Disclosures: Philippe Lachapelle reports speaker honoraria from AstraZeneca, Boehringer Ingelheim, GlaxoSmithKline, Novartis and Sanofi-Regeneron. Simon Couillard reports the following: non-restricted research grants from AstraZeneca, bioMérieux, Sanofi-Genzyme-Regeneron, the Fondation Québécoise en Santé Respiratoire, the NIHR Oxford BRC and the Quebec Respiratory Health Research Network; speaker honoraria from AstraZeneca, GlaxoSmithKline, Sanofi-Regeneron and Valeo Pharma; consultancy fees for AstraZeneca, FirstThought, GlaxoSmithKline and Sanofi-Regeneron; sponsorship to attend/speak at international scientific meetings by/for AstraZeneca and Sanofi-Regeneron; is an advisory board member and will have stock options for Biometry Inc, a company which is developing a FeNO device (myBiometry); advised the Institut national d'excellence en santé et services sociaux (INESSS) for an update of the asthma general practice information booklet for general practitioners; is the holder of the Association Pulmonaire du Québec's Research Chair in Respiratory medicine and is a clinical research scholar of the Fonds de recherche du Québec. Laurence Désy has no conflicts of interest to declare in relation to this article.

Acknowledgements: The authors wish to thank the patients with severe asthma and the respiratory therapists, nursing team and pulmonologists at the Centre Hospitalier Universitaire de Sherbrooke for their inspiration and support.

Review process: Double-blind peer review.

Compliance with ethics: This article is an opinion piece and does not report on new clinical data, or any studies with human or animal subjects performed by any of the authors.

Data availability: Data sharing is not applicable to this article as no datasets were generated or analyzed during the writing of this article.

Authorship: The named authors meet the International Committee of Medical Journal Editors (ICMJE) criteria for authorship of this manuscript, take responsibility for the integrity of the work as a whole, and have given final approval for the version to be published.

Access: This article is freely accessible at touchRESPIRATORY.com © Touch Medical Media 2023

Received: 15 June 2023

Accepted: 21 July 2023

Published online: 10 October 2023

Citation: touchREVIEWS in Respiratory & Pulmonary Diseases. 2024;9(1):Online ahead of journal publication

Corresponding author: Simon Couillard, Centre Hospitalier Universitaire de Sherbrooke, 3001, 12e Avenue Nord, pièce 2616, Sherbrooke, Québec, Canada. E: s.couillard@usherbrooke.ca

Support: Funding from the research chair in Respiratory Health of the Lung Association of Quebec and the Fonds de recherche du Québec.

The topic of the carbon footprint of human activities is one frequently discussed worldwide. There have been conscious efforts across multiple fields, including healthcare, to reduce their carbon footprint. In Canada, 4.6% of greenhouse gases (GHG) are manufactured by the healthcare institutions, ranking it second worldwide.^{1,2} The treatments for asthma and chronic obstructive pulmonary disease (COPD) are two of the main contributors, this includes pressurized metered dose inhalers (pMDIs), dry powder inhalers (DPIs) and soft mist inhalers (SMIs). These highly prevalent diseases affect 11% and 10% of the Western populations, respectively.³ This editorial aims to highlight the impact of pressurized metered dose inhalers on the environment, discussing alternatives and changes to current diagnostic and therapeutic practices, public awareness, potential changes to regulations and pathways moving forward.


Impacts of pressurized metered dose inhalers

Used in patients with respiratory diseases like asthma and COPD, pMDIs represent 0.03% of the annual worldwide carbon footprint.⁴ Due to their effects on the environment, the Montreal Protocol^{5,6} was established in 1987 and signed by 197 countries, regulating the use of many substances affecting the ozone layer, such as the chlorofluorocarbons (CFCs) contained in pMDIs. Following this historical pact, CFCs were replaced by hydrofluorocarbons (HFCs), which had potentially fewer consequences on global warming. However, HFCs remain important vectors for GHG emissions.⁶ HFCs are synthetic gases used as a substitute for ozone-depleting substances, however, they do still contribute to GHG emissions.⁷ For instance, a single dose from a pMDI, which includes two inhalations, is the equivalent of driving two kilometres with a hybrid vehicle, with regards to carbon dioxide emissions.⁸

Alternatives to pressurized metered dose inhalers using hydrofluorocarbons

Beyond pMDIs, different types of inhalers with different delivery mechanisms are available on the market. DPIs use peak inspiratory flow to deliver the medication inside the respiratory tract without the use of propellants. Comparing the different inhaler types on the market, one can notice that DPIs and SMIs have a carbon footprint approximately 10 times smaller, compared with pMDIs.⁸ However, >70% of inhalers currently sold to adult patients are pMDIs.⁹ It must be mentioned that, if the inhalation technique/strength with DPIs is adequate, their efficacy parallels that of pMDIs, but has a much lower impact on the environment.¹⁰ In a recent study, patients using pMDIs were switched to DPIs, resulting in reducing

Figure 1: The excessive carbon footprint of inhaler devices used in airway disease and its remedies



COPD = chronic obstructive pulmonary disease; GHG = greenhouse gases.

their carbon footprint by more than half with no impact on their disease control.¹¹ If 5% of patients per year substituted their pMDIs to DPIs, there would be a reduction of 27% of GHGs related to inhaler use over the course of 10 years.¹²

There are many factors to consider when prescribing inhalers. For example, the therapeutic efficacy of DPIs will be lower for individuals with advanced pulmonary disease, due to inadequate drug delivery. Indeed, with inspiratory muscle weakness and associated pulmonary hyperinflation, the maximal inspiratory pressure is insufficient for dry powder inhalation.¹³ Additionally, it can be difficult for patients to inhale properly during an acute airway attack. Personal preference and the varying costs of inhalers must also be considered. Nevertheless, in a study comparing costs, device handling and the environmental impact of the inhalers, only 14% of the participants indicated that carbon footprint was not a decisional factor in the choice of inhalers.¹⁴

The importance of an established diagnosis for prescribing or renewing treatment

A formal diagnosis of asthma or COPD with objective tests is crucial. Unfortunately, in one study, it was found that 33% of randomly recruited patients that received a diagnosis for asthma within the last 5 years had no evidence of the disease during the pulmonary function test.¹⁵ Of those individuals in whom an asthma diagnosis was ruled out, 79% had used inhalers prior to the re-evaluation, and thus have contributed to the ecological damages caused by HFCS.¹⁵

The frequently overlooked inhaler technique

Aside from considering a substitution from pMDIs to DPIs and/or SMIs, it is crucial to educate patients in proper inhaler technique, since most GHG emissions are produced during the inhalation phase.⁷ When choosing a pMDI, the use of a spacer allows for the optimal medication quantity without wasting and overutilization. This also reduces the number of

inhalers used yearly, as less drug is used with each use of the inhaler, which decreases GHG. An estimated 68% of patients do not properly use their inhalers, perhaps due to most of them not receiving adequate education on the proper techniques.¹⁶ Furthermore, for many patients using pMDIs, it is often challenging to determine if the inhaler is empty, with the absence of a numerical dial indicating the number of remaining doses. In contrast, a dial is present on the majority of the DPI devices. Studies have shown that 40% of patients use empty inhalers, which contributes to poor control of the disease.¹⁷ Conversely, some patients tend to renew their inhalers prior to finishing all of the doses, which increases product waste and overutilization. In summary, to reduce GHG emissions in airway disease, it is important to educate patients on the use of inhalers, identify poor symptom control and discuss strategies to prevent disease exacerbations.

Controlling asthma to control greenhouse gas emissions

The impact of inadequate asthma control must not be underestimated, both in terms of morbidity and carbon footprint. Compared with patients with adequate control of their disease, poorly controlled airway disease is associated with threefold greater GHG emissions.¹⁸ The most adequate way to maintain a proper control of disease is to limit tobacco use, identify and target unchecked airway inflammation and reduce exposure to allergens, but also to verify treatment use and inhalation technique with patients. Furthermore, in patients aged 12 and up with well-controlled mild asthma, guidelines from the Global Initiative for Asthma recommend that treatment with budesonide-formoterol inhalers (usually a DPI) is preferred to a daily inhaled corticosteroid paired with a short-acting bronchodilator (usually pMDIs).¹⁹ Airway disease that is well-controlled requires less reliever use and healthcare use, effectively reducing the associated carbon footprint.

Returning inhalers to the local pharmacist

The carbon footprint associated with the manufacturing and disposal of pMDIs may represent an estimated 30% of the total footprint of the inhaler.⁹ The incineration and recycling of the devices reduces the impact on global warming.⁷ To this day, no systematic programme has been implemented with regards to recycling or packaging of the inhalers. At the very least, devices should be disposed of adequately.²⁰

Awareness

There are programmes in place to educate patients and healthcare professionals in taking an eco-friendly approach. CASCADE is an organization funded by the Canadian Government partnering with the Canadian Thoracic Society, which is similar to its counterpart PrescQIPP in the UK.^{21,22} These programmes aim to guide the healthcare systems towards a more durable environment with a neutral carbon use. To reach their objectives, they offer multiple resources explaining how to reduce carbon emissions. These include videos on inhalation techniques, information posters and letters explaining the proper use of inhalers.²³

Regulations

Regulation can limit the use of polluting gases as witnessed in the Montreal Protocol.⁵ In addition to the efforts put in since 1987, the Kigali

Amendment to the Montreal Protocol in 2019 aims to reduce carbon emissions further, imposing a mandated reduction of HFCs worldwide.⁶ Hydrofluoroolefins (HFOs) are currently in development to assure an alternative pMDIs propellant.²⁴ Replacing HFCs with HFOs would substantially reduce the global warming potential of pMDIs.

Conclusions

In conclusion, healthcare systems must take part in reducing their carbon footprint. This is especially true for airway disease and its associated inhalers (*Figure 1*). pMDIs constitute a healthcare risk by decreasing the air quality and as a result increase symptoms and exacerbations of pulmonary diseases. The choice of inhalers depends on a multitude of factors, including personal preference, price, inhalation technique and simplicity, as well as their carbon footprint. Efforts have been made to reduce GHG emissions, such as the CASCADE program, recycling programmes in pharmacies, the Kigali Amendment aiming to progressively reduce the HFCs, and developing new alternatives to the propulsion mechanism with HFO.

Finally, a proposition: after CFCs, why not regulate the progressive removal of HFC-based pMDIs in order to promote HFO-based pMDIs? We can and should be as ambitious for the environment as for our patients.



- Romanello M, McGushin A, Di Napoli C, et al. The 2021 report of the Lancet Countdown on health and climate change: Code red for a healthy future. *Lancet*. 2021;398:1619–62. DOI: 10.1016/S0140-6736(21)01787-6.
- Eckelman MJ, Sherman JD, MacNeill AJ. Life cycle environmental emissions and health damages from the Canadian healthcare system: An economic-environmental-epidemiological analysis. *PLoS Med*. 2018;15:e1002623. DOI: 10.1371/journal.pmed.1002623.
- Public Health Agency of Canada. Report from the Canadian Chronic Disease Surveillance System: Asthma and chronic obstructive pulmonary disease (COPD) in Canada, 2018. 2018. Available at: www.canada.ca/content/dam/phac-aspc/documents/services/publications/diseases-conditions/asthma-chronic-obstructive-pulmonary-disease-canada-2018/pub-eng.pdf (Date last accessed: 30 May 2023).
- Association pulmonaire du Québec. Crise climatique. Available at: <https://poumonquebec.ca/sante-pulmonaire/environnement/crise-climatique/> (Date last accessed: 15 June 2023). French.
- The U.S. Department of State. The Montreal protocol on substances that deplete the ozone layer. Available at: www.state.gov/key-topics-office-of-environmental-quality-and-transboundary-issues/the-montreal-protocol-on-substances-that-deplete-the-ozone-layer/ (Date last accessed: 13 September 2023).
- UN Environment Programme. About Montreal protocol. Available at: www.unep.org/ozonaction/who-we-are/about-montreal-protocol (Date last accessed: 15 June 2023).
- Wilkinson AJK, Braggins R, Steinbach I, Smith J. Costs of switching to low global warming potential inhalers. An economic and carbon footprint analysis of NHS prescription data in England. *BMJ Open*. 2019;9:e028763. DOI: 10.1136/bmjopen-2018-028763.
- Panigone S, Sandri F, Ferri R, et al. Environmental impact of inhalers for respiratory diseases: Decreasing the carbon footprint while preserving patient-tailored treatment. *BMJ Open Respir Res*. 2020;7:e000571. DOI: 10.1136/bmjresp-2020-000571.
- Janson C, Henderson R, Löfdahl M, et al. Carbon footprint impact of the choice of inhalers for asthma and COPD. *Thorax*. 2020;75:82–4. DOI: 10.1136/thoraxjnl-2019-213744.
- Brocklebank D, Ram F, Wright J, et al. Comparison of the effectiveness of inhaler devices in asthma and chronic obstructive airways disease: A systematic review of the literature. *Health Technol Assess*. 2001;5:1–149. DOI: 10.3310/hta5260.
- Woodcock A, Janson C, Rees J, et al. Effects of switching from a metered dose Inhaler to a dry powder Inhaler on climate emissions and asthma control: Post-hoc analysis. *Thorax*. 2022;77:1187–92. DOI: 10.1136/thoraxjnl-2021-218088.
- Perrigotti D, Storham C, Panigone S, et al. Reducing carbon footprint of inhalers: Analysis of climate and clinical implications of different scenarios in five European countries. *BMJ Open Respir Res*. 2021;8:e001071. DOI: 10.1136/bmjresp-2021-001071.
- Clark AR, Weers JG, Dhand R. The confusing world of dry powder Inhalers: It is all about Inspiratory pressures, not Inspiratory flow rates. *J Aerosol Med Pulm Drug Deliv*. 2020;33:1–11. DOI: 10.1089/jamp.2019.1556.
- Liew K, Wilkinson A. P280 how do we choose Inhalers? patient and physician perspectives on environmental, financial and ease-of-use factors. *Thorax*. 2017;72:A235–7. DOI: 10.1136/thoraxjnl-2017-210983.422.
- Aaron SD, Vandemheen KL, FitzGerald JM, et al. Reevaluation of diagnosis in adults with physician-diagnosed asthma. *JAMA*. 2017;317:269–79. DOI: 10.1001/jama.2016.19627.
- Haidl P, Heindl S, Siemon K, et al. Inhalation device requirements for patients' inhalation maneuvers. *Respir Med*. 2016;118:65–75. DOI: 10.1016/j.rmed.2016.07.013.
- Conner JB, Buck PO. Improving asthma management: The case for mandatory inclusion of dose counters on all rescue Bronchodilators. *J Asthma*. 2013;50:658–63. DOI: 10.3109/02770903.2013.789056.
- Wilkinson A, Maslova E, Janson C, et al. Greenhouse gas emissions associated with asthma care in the UK: Results from SABINA CARBON. *Eur Respir J*. 2022;58:A76. DOI: 10.1183/13993003.congress-2021.OA76.
- Global Initiative for Asthma. 2023 GINA Report, Global Strategy for Asthma management and prevention. 2023. Available at: <https://ginasthma.org/2023-gina-main-report/> (Date last accessed: 16 August 2023).
- Health Products Stewardship Association. Returning Medications. Available at: <https://healthsteward.ca/consumers/returning-medications/> (Date last accessed: 15 June 2023).
- CASCADES. Available at: <https://cascadescanada.ca/> (Date last accessed: 13 September 2023).
- PrescQIPP. Available at: <https://prescqipp.info/> (Date last accessed: 13 September 2023).
- Cascades Canada. Sustainable pharmacy and prescribing: Implementation resources for sustainable pharmacy and prescribing practices. Available at: <https://cascadescanada.ca/action-areas/pharmacy-and-prescribing/> (Date last accessed: 15 June 2023).
- Hargreaves C, Budgen N, Whiting A, et al. S60 A new medical Propellant HFO-1234Ze(E): Reducing the environmental impact of inhaled medicines. *Thorax*. 2022;77:A38–9. DOI: 10.1136/thorax-2022-BTSabstracts.66.